

FORM PTO-1390
(REV 10-95)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. §371**

UMARY 5

U.S. APPLICATION NO. (If known, see 37 CFR §1.5)

09/856044

INTERNATIONAL APPLICATION NO.

PCT/US99/27191

INTERNATIONAL FILING DATE

17 NOVEMBER 1999

PRIORITY DATE CLAIMED

17 NOVEMBER 1998

TITLE OF INVENTION

METHODS FOR IDENTIFYING AND CLASSIFYING ORGANISMS BY MASS SPECTROMETRY AND DATABASE SEARCHING

APPLICANT(S) FOR DO/EO/US



DEMIREV, Plamen, A., et al.

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. §371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. §371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. §371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. §371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. §371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. §371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. §371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. §371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. §371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. §371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 C.F.R. §§1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 C.F.R. §§3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.
 - ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☐ Other items or information:

U.S. APPLICATION NO. (if known, see 37 CFR §1.5) <div style="font-size: 2em; font-weight: bold; margin-top: 5px;">09/856044</div>		INTERNATIONAL APPLICATION NO. PCT/US99/27191		ATTORNEY'S DOCKET NUMBER UMARY 5 WO	
17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR §1.492 (a) (1) - (5)): Search Report has been prepared by the EPO or JPO..... \$860.00 International preliminary examination fee paid to USPTO (37 CFR §1.482)..... \$690.00 No international preliminary examination fee paid to USPTO (37 CFR §1.482) but international search fee paid to USPTO (37 CFR §1.445(a)(2))..... \$710.00 Neither international preliminary examination fee (37 CFR §1.482) nor international search fee (37 CFR §1.445(a)(2)) paid to USPTO..... \$1000.00 International preliminary examination fee paid to USPTO (37 CFR §1.482) and all claims satisfied provisions of PCT Article 33(2)-(4)..... \$100.00 <div style="text-align: right; margin-top: 5px;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>				CALCULATIONS PTO USE ONLY <div style="border: 1px solid black; height: 100px; margin-top: 5px;"></div>	
Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 C.F.R. §1.492(e)). <input type="checkbox"/> 20 <input type="checkbox"/> 30					
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	11 - 20 =	0	x \$ 18.00	\$0.00	
Independent claims	1 - 3 =	0	x \$ 80.00	\$0.00	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)				+ \$ 270.00	
TOTAL OF ABOVE CALCULATIONS =				\$860.00	
Reduction of 1/2 for filing by small entity, if applicable. Applicant qualifies.				\$430.00	
SUBTOTAL =				\$430.00	
Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 C.F.R. §1.492(f)). <input type="checkbox"/> 20 <input type="checkbox"/> 30					
TOTAL NATIONAL FEE =				\$430.00	
Fee for recording the enclosed assignment (37 C.F.R. §1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 C.F.R. §§3.28, 3.31). \$40.00 per property.					
TOTAL FEES ENCLOSED =				\$430.00	
				Amount to be refunded:	
				charged:	
a. <input checked="" type="checkbox"/> A check in the amount of <u>\$430.00</u> to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. <u>13-3402</u> in the amount of \$_____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>13-3402</u> . A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 C.F.R. §§1.494 or 1.495 has not been met, a petition to revive (37 C.F.R. §1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO: Customer Number 23,599 <div style="display: flex; justify-content: space-between; align-items: center; margin-top: 10px;"> <div style="text-align: center;">  <div style="font-size: 1.5em; font-weight: bold;">23599</div> <div style="font-size: 0.8em;">PATENT TRADEMARK OFFICE</div> </div> <div style="text-align: right; width: 60%;"> <div style="font-size: 1.5em; margin-bottom: 5px;">  </div> <div style="border-top: 1px solid black; width: 100%;"></div> <div style="font-size: 0.9em;">SIGNATURE</div> <div style="border-top: 1px solid black; width: 100%;"></div> <div style="font-size: 0.9em;">NAME</div> <div style="border-top: 1px solid black; width: 100%;"></div> <div style="font-size: 0.9em;">REGISTRATION NUMBER</div> </div> </div>					
Filed: 17 MAY 2001 RJT:kms					

IN THE UNITED STATES DESIGNATED/ELECTED OFFICE

International Application No. : PCT/US99/27191
International Filing Date : 17 NOVEMBER 1999
Priority Date(s) Claimed : 17 NOVEMBER 1998
Applicant(s) (DO/EO/US) : DEMIREV, Plamen, et al.
Title: METHODS FOR IDENTIFYING AND CLASSIFYING ORGANISMS BY MASS
SPECTROMETRY AND DATABASE SEARCHING

PRELIMINARY AMENDMENT

Commissioner for Patents
Washington, D.C. 20231

SIR:

Although the claims of the above application were amended under Article 34 during the International Phase, applicants respectfully request that examination be based on the claims as originally filed.

Prior to calculating the national fee, and prior to examination in the National Phase of the above-identified International application, please amend as follows:

IN THE SPECIFICATION:


Page 1, before the first paragraph, please insert the following paragraph:

--This application claims the benefit of priority of U.S. Provisional Applications 60/108,696 filed November 17, 1998 and 60/120,679 filed February 19, 1999.--

REMARKS

The purpose of this Preliminary Amendment is to request examination of the application as filed and to insert a claim of priority. Applicants reserve the right to reintroduce claims to canceled combined subject matter.

Respectfully submitted,



Richard J. Traverso, Registration No. 30,595
Attorney for Applicants
MILLEN, WHITE, ZELANO & BRANIGAN, P.C.
Arlington Courthouse Plaza 1
2200 Clarendon Boulevard, Suite 1400
Arlington, VA 22201
Direct Dial: 703-812-5310
Facsimile: 703-243-6410
Email: traverso@mwzb.com

RJT:jmm

Filed: 17 MAY 2001

A new paragraph was inserted in the specification, therefore no marked up version is necessary.

Cognitive Function		Psychiatric Function		Quality of Life	
Measure	Score	Measure	Score	Measure	Score
MMSE	24.5	Beck Depression Inventory	15.2	WHOQOL-BREF	78.5
MoCA	26.8	Beck Anxiety Inventory	12.1	WHOQOL-REF	82.3
Trail Making Test	18.2	Stress Management Scale	85.4	WHOQOL-REF	85.1
Digit Span	12.5	Life Satisfaction Scale	72.3	WHOQOL-REF	88.9
Block Design	22.1	Life Satisfaction Scale	75.6	WHOQOL-REF	91.2
Verbal Fluency	15.3	Life Satisfaction Scale	78.9	WHOQOL-REF	93.5
Stroop Test	19.7	Life Satisfaction Scale	81.2	WHOQOL-REF	95.8
Digit Span	13.1	Life Satisfaction Scale	83.5	WHOQOL-REF	98.1
Block Design	23.4	Life Satisfaction Scale	85.8	WHOQOL-REF	100.0
Verbal Fluency	16.7	Life Satisfaction Scale	88.1	WHOQOL-REF	102.3
Stroop Test	20.9	Life Satisfaction Scale	90.4	WHOQOL-REF	104.6
Digit Span	13.8	Life Satisfaction Scale	92.7	WHOQOL-REF	106.9
Block Design	24.7	Life Satisfaction Scale	95.0	WHOQOL-REF	109.2
Verbal Fluency	17.9	Life Satisfaction Scale	97.3	WHOQOL-REF	111.5
Stroop Test	22.1	Life Satisfaction Scale	99.6	WHOQOL-REF	113.8
Digit Span	14.5	Life Satisfaction Scale	101.9	WHOQOL-REF	116.1
Block Design	25.8	Life Satisfaction Scale	104.2	WHOQOL-REF	118.4
Verbal Fluency	19.1	Life Satisfaction Scale	106.5	WHOQOL-REF	120.7
Stroop Test	23.4	Life Satisfaction Scale	108.8	WHOQOL-REF	123.0
Digit Span	15.2	Life Satisfaction Scale	111.1	WHOQOL-REF	125.3
Block Design	27.1	Life Satisfaction Scale	113.4	WHOQOL-REF	127.6
Verbal Fluency	20.3	Life Satisfaction Scale	115.7	WHOQOL-REF	129.9
Stroop Test	24.7	Life Satisfaction Scale	118.0	WHOQOL-REF	132.2
Digit Span	16.5	Life Satisfaction Scale	120.3	WHOQOL-REF	134.5
Block Design	28.4	Life Satisfaction Scale	122.6	WHOQOL-REF	136.8
Verbal Fluency	21.6	Life Satisfaction Scale	124.9	WHOQOL-REF	139.1
Stroop Test	26.0	Life Satisfaction Scale	127.2	WHOQOL-REF	141.4
Digit Span	17.8	Life Satisfaction Scale	129.5	WHOQOL-REF	143.7
Block Design	29.7	Life Satisfaction Scale	131.8	WHOQOL-REF	146.0
Verbal Fluency	22.9	Life Satisfaction Scale	134.1	WHOQOL-REF	148.3
Stroop Test	27.1	Life Satisfaction Scale	136.4	WHOQOL-REF	150.6
Digit Span	19.0	Life Satisfaction Scale	138.7	WHOQOL-REF	152.9
Block Design	30.4	Life Satisfaction Scale	140.7	WHOQOL-REF	155.2
Verbal Fluency	24.0	Life Satisfaction Scale	143.0	WHOQOL-REF	157.5
Stroop Test	28.4	Life Satisfaction Scale	145.3	WHOQOL-REF	159.8
Digit Span	20.2	Life Satisfaction Scale	147.6	WHOQOL-REF	162.1
Block Design	31.7	Life Satisfaction Scale	149.6	WHOQOL-REF	164.4
Verbal Fluency	25.1	Life Satisfaction Scale	151.9	WHOQOL-REF	166.7
Stroop Test	29.7	Life Satisfaction Scale	154.2	WHOQOL-REF	169.0
Digit Span	21.4	Life Satisfaction Scale	156.5	WHOQOL-REF	171.3
Block Design	33.0	Life Satisfaction Scale	158.5	WHOQOL-REF	173.6
Verbal Fluency	26.2	Life Satisfaction Scale	160.8	WHOQOL-REF	175.9
Stroop Test	30.0	Life Satisfaction Scale	163.1	WHOQOL-REF	178.2
Digit Span	22.6	Life Satisfaction Scale	165.4	WHOQOL-REF	180.5
Block Design	34.3	Life Satisfaction Scale	167.4	WHOQOL-REF	182.8
Verbal Fluency	27.1	Life Satisfaction Scale	169.7	WHOQOL-REF	185.1
Stroop Test	31.3	Life Satisfaction Scale	172.0	WHOQOL-REF	187.4
Digit Span	23.8	Life Satisfaction Scale	174.3	WHOQOL-REF	189.7
Block Design	35.6	Life Satisfaction Scale	176.3	WHOQOL-REF	192.0
Verbal Fluency	28.0	Life Satisfaction Scale	178.6	WHOQOL-REF	194.3
Stroop Test	32.6	Life Satisfaction Scale	180.9	WHOQOL-REF	196.6
Digit Span	25.0	Life Satisfaction Scale	183.2	WHOQOL-REF	198.9
Block Design	36.9	Life Satisfaction Scale	185.2	WHOQOL-REF	201.2
Verbal Fluency	29.1	Life Satisfaction Scale	187.5	WHOQOL-REF	203.5
Stroop Test	33.9	Life Satisfaction Scale	189.8	WHOQOL-REF	205.8
Digit Span	26.2	Life Satisfaction Scale	192.1	WHOQOL-REF	208.1
Block Design	38.2	Life Satisfaction Scale	194.1	WHOQOL-REF	210.4
Verbal Fluency	30.2	Life Satisfaction Scale	196.4	WHOQOL-REF	212.7
Stroop Test	34.2	Life Satisfaction Scale	198.7	WHOQOL-REF	215.0

PTO/PCT Rec'd 17 MAY 2001

**METHODS FOR IDENTIFYING AND CLASSIFYING ORGANISMS BY
MASS SPECTROMETRY AND DATABASE SEARCHING**

Cross-reference to Applications

5 This application claims the benefit of U.S. Provisional Application Nos.
60/108,696, filed November 17, 1998, and 60/120,679, filed February 19, 1999, which
are hereby incorporated by reference in their entirety.

Background of the Invention

10 The development of methods of rapidly identifying and characterizing biological
materials, such as microorganisms and cells, is major focus of academic and industrial
research. The need for such methods has been felt in the health, laboratory, and
environmental industries. In medicine, for example, the exponential rise in antibiotic-
resistant bacteria and emerging viral disease has caused a crisis in the health-care and
food industries. As a result, there has been a continued pressure to find new, reliable, and
rapid means of characterizing pathological and disease-causing organisms. Similarly, the
15 threats of biological warfare and terrorist activities which have been felt world-wide has
caused an escalated search for ways of identifying putative biological agents, especially
in the field, in airports, and in other public areas.

20 Coupled with the need for advanced biological agent detection methods has been
an escalating effort in the sequencing of DNA from all types of organisms and
identifying expressed genes. The complete genomic sequences of a number of
microorganisms had been completed. The availability of such information about genome
and proteome of whole organisms is an important reservoir to be exploited for identifying
and characterizing unknown and known organisms.

Description of the Drawings

Fig. 1: Molecular mass distribution (in bins of 1 kDa) of proteins deposited in the SwissPROT/TrEMBL sequence database: **a)** all prokaryotic proteins, and **b)** all proteins from *B.subtilis*.

5 **Fig. 2:** Positive ion MALDI spectra from: **a)** *B.subtilis* (8 hours growth time), matrix - SA; and **b)** *E.coli* (32 hours growth time), matrix - CHCA.

Fig. 3: Positive ion MALDI spectra from: **a)** *E.coli* (32 hours growth time), matrix - MCA:SA mixture; and **b)** *E.coli* (8 hours growth time), matrix - CHCA.

10 **Fig. 4:** Number of proteins combined from *B.subtilis* and *E.coli* with masses within a predetermined mass window (in ppm) as a function of molecular mass.

Fig. 5: Positive ion MALDI spectrum from a mixture of *B.subtilis* and *E.coli*, matrix - SA.

Fig. 6: An example of a flow chart for microorganism and cell identification by mass spectrometry and database searching.

15 **Description of the Invention**

The present invention relates to compositions of matter, instruments, processes (e.g., as carried using computer software and/or hardware), and methods, for identifying, classifying, and/or characterizing biological materials by measuring the molecular weights of protein constituents in such materials and using the molecular information to
20 deduce the organismic source of the biological materials. In preferred embodiments of the invention, biological materials comprising proteins can be subjected to mass spectrometry, or other suitable means for determining mass, in order to determine the molecular weights of the protein constituents. The resulting molecular weight information of the protein constituents can then be used to query databases which
25 contain, among other information, lists of protein molecular weight information and the identity of the organism source from which the information was derived. By comparing the set of protein molecular masses of an unknown, as determined, for instance, in a mass spectrum, against a database containing the molecular masses of proteins present in known organisms, the unknown can be rapidly and reliably identified, classified, or
30 characterized.

-3-

5 The present invention presents a method for rapid identification, classification, and/or characterization of biological materials, such as microorganisms, organisms, organs, tissues, cells, subcellular materials, and the like, which exploits the wealth of information contained in genome and protein sequence databases. The massive efforts to sequence the human genome has brought about a rapid increase in the speed with which DNA sequences from all species are being accumulated in publicly available computer databases. As a result, the complete genomes of many different organisms are now completely known (e.g., National Center for Biotechnology Information (NIH), <http://www.ncbi.nlm.nih.gov/Entrez/Genome/org.html>; The *C. elegans* Sequencing Consortium, *Science* **1998**, 282, 2012–2018; TIGR Microbial Database, <http://www.tigr.org/tdb/mdb/mdb.html>). See, also, Table 6. There exists complementarity between the genome of an organism, and its respective proteome, i.e. the dynamic entity set of all expressed proteins. In databases, such complementarity is realized via assignment of an amino acid sequence to each “open reading frame” (ORF) in a DNA sequence. By using bioinformatics tools, the complete proteomes of organisms with established DNA sequences have been made available and accessible, e.g., through the Internet. Characterization of such organisms can be achieved through knowledge of their complete genomes or complementary proteomes.

20 In accordance with the present invention, any instrument, method, process, etc. can be utilized to determine the molecular weight of proteins in a sample. A preferred method of obtaining molecular weight is by mass spectrometry, where protein molecules in a sample are ionized and then the resultant mass and charge of the protein ions are detected and determined.

25 Any suitable instrument, method, process, etc. for carrying out mass spectroscopy can be utilized. To use mass spectrometry to analyze proteins, it is preferred that the protein be converted to a gas-ion phase. Various methods of protein ionization are useful, including, e.g., fast ion bombardment (FAB), plasma desorption, laser desorption, thermal desorption, preferably, electrospray ionization (ESI) and matrix-assisted laser desorption/ionization (MALDI). Many different mass analyzers are available for peptide and protein analysis, including, but not limited to, Time-of-Flight (TOF), ion trap (ITMS), Fourier transform ion cyclotron (FTMS), quadrupole ion trap, and sector

-4-

(electric and/or magnetic) spectrometers. See, e.g., U.S. Pat. No. 5,572,025 for an ion-trap MS.

Mass analyzers can be used alone, or in combination to form tandem mass spectrometers. In the latter case, a first mass analyzer can be use to separate the protein ions (precursor ion) from each other and determine the molecular weights of the various protein constituents in the sample. A second mass analyzer can be used to analyze the separated constituents, e.g., by fragmenting the precursor ions into product ions. Any desired combination of mass analyzers can be used, including, e.g., triple quadrupoles, tandem time-of-flights, ion traps, and/or combinations thereof.

Different kinds of detectors can be used to detect the protein ions. For example, destructive detectors can be utilized, such as ion electron multipliers or cryogenic detectors (e.g., U.S. Pat. No. 5,640,010). Additionally, non-destructive detectors can be used, such as an ion trap which is utilized in an ion current pick-up devices which are utilized in quadrupole ion trap mass analyzers or FTMS.

Any source of proteins can be used in accordance with the present invention, including whole organisms, such as multicellular and unicellular organisms, organs, tissues, cells, subcellular structures, and mixtures thereof. Various microorganisms can be utilized, e.g., archeabacteria, bacteria, chlamydiae, rickettsia, viruses, mycoplasma, molds, yeasts, protozoa, algae, prions, etc. Cells, microorganisms, etc. can be genetically-engineered, altered, modified, etc. Proteins can be extracted from intact or treated materials. Any substrate comprising a biological material can be used. For instance, it may be desirable to characterize organisms found on surfaces, in food, in biological fluids, such as saliva, urine, fecal matter, blood, lymph, or plasma, on materials used to wipe surfaces suspected of containing organisms, in hair, objects handled or contacted by organisms, etc.

Any method of preparing samples for analysis can be used. For MALDI-TOF, a number of sample preparation methods can be utilized including, dried droplet (Karas and Hillenkamp, *Anal. Chem.*, 60:2299-2301, 1988), vacuum-drying (Winberger et al., *In Proceedings of the 41st ASMS Conference on Mass Spectrometry and Allied Topics*, San Francisco, May 31-June 4, 1993, pp. 775a-b), crush crystals (Xiang et al., *Rapid Comm. Mass Spectrom.*, 8:199-204, 1994), slow crystal growing (Xiang et al., *Org. Mass Spectrom.*, 28:1424-1429, 1993); active film (Mock et al., *Rapid Comm. Mass Spectrom.*,

6:233-238, 1992; Bai et al., *Anal. Chem.*, 66:3423-3430, 1994), pneumatic spray (Kochling et al., *Proceedings of the 43rd ASMS Conference on Mass Spectrometry and Allied Topics*; Atlanta, GA, May 21-26, 1995, p1225); electrospray (Hensel et al., *Proceedings of the 43rd ASMS Conference on Mass Spectrometry and Allied Topics*; Atlanta, GA, May 21-26, 1995, p947); fast solvent evaporation (Vorm et al., *Anal. Chem.*, 66:3281-3287, 1994); sandwich (Li et al., *J. Am. Chem. Soc.*, 118:11662-11663, 1996); and two-layer methods (Dai et al., *Anal. Chem.*, 71:1087-1091, 1999). See also, e.g., Liang et al., *Rapid Commun. Mass Spectrom.*, 10:1219-1226, 1996; van Adrichem et al., *Anal. Chem.*, 70:923-930, 1998. For example, samples of microorganisms can be lyophilized, extracted into a solution, such as a 70:30 solution of CH₃CN:0.1% trifluoroacetic acid, and then embedded in the matrix. Various matrices can be used, e.g., sinapinic acid, 2,5-dihydroxybenzoic acid, alpha-cyano-4-hydroxycinnamic acid. A sample can be processed in various ways prior to addition to the matrix. For instance, the sample can be extracted, subjected to corona discharge, chromatography, such as HPLC, etc., e.g., to remove particular unwanted constituents (such as lipids, small molecules, high molecular weight constituents) before mass spectrometry.

MALDI-TOF can detect proteins at the attamole level over a wide range of molecular weights. Masses can be determined, e.g., as low as 1000, and as high as a hundred-thousand daltons. Any range of molecular weights can be used in accordance with the present invention, e.g., about 4000-20,000. Masses accurate to about 50 ppm or better will be most reliable in the general case.

In some cases, it may be desirable to obtain more information on a particular protein identified in a mass spectrum. One instance is where a specific peak matches more than one different protein in the searched databases. This can happen when different proteins share the same, or similar, molecular weights. A search of a database in such a case might reveal more than one protein which corresponds to the measured molecular weight of a protein in the sample. To determine which protein in the database corresponds to the peak at issue, the peak protein can be separated from the mixture and subjected to further physical characterization. Separation can be accomplished by any suitable method, e.g., conventional techniques involving, e.g., by cell lysis, extraction, and two-dimensional gel chromatography, capillary electrophoresis, or high performance liquid chromatography

-6-

Useful information includes, amino acid composition, amino acid sequence, proteolytic and enzymatic cleavage patterns, isoelectric point, hydrophobicity, and other physical characteristics. Another scenario where additional information on a protein in spectrum may be warranted is where such protein has not matched any of the proteins listed in the database. Thus, such information can be useful to increase the specificity of the approach.

As mentioned, characterization of individual proteins in mixture can be accomplished using any suitable means. The expanding requirements in proteomics, e.g. for rapid identification of proteins present in a mixture in picomolar amounts, have resulted in the development of powerful MS-based procedures for identity assignment of individual proteins. [9-14] These methods include, but are not limited to, chemical/enzymatic digestion of material (obtained from a single spot in a two-dimensional gel electropherogram, or by other suitable chromatographic technique) and mass spectral determination of the molecular masses of the protein and resulting peptides (peptide mapping). Making use of already available information in protein sequence databases, a comparison can be made between proteolytic peptide mass patterns generated "in silico," and experimentally-observed peptide masses. A "hit-list" can be compiled, ranking candidate proteins in the database, based on (among other criteria) number of matches between the proteolytic fragments. Several Web sites are accessible that provide software for protein identification on-line, based on peptide mapping and sequence database search strategies.

[15] Methods of peptide mapping and sequencing using MS are described in WO95/25281, U.S. Pat. No. 5,538,897, U.S. Pat. No. 5,869,240, U.S. Pat. No. 5,572,025, U.S. Pat. No. 5,696,376. See, also, Yates, J. Mass Spec., 33:1-19. 1998. The present invention can also be combined with methods that detect small molecules (other than proteins) in samples, such as the method described in WO98/09314. The latter method only measure molecules in the range of about 500-1500 Da, and not more than 1876 Da.

Data collected from a mass spectrometer typically comprises the intensity and mass to charge ratio for each detected event. Spectral data can be recorded in any suitable form, including, e.g., in graphical, numerical, or electronic formats, either in digital or analog form. Spectra is preferably recorded in a storage medium, including,

-7-

e.g., magnetic, such as floppy disk, tape, or hard disk; optical, such as CD-ROM or laser-disc; or, ROM-CHIPS.

5 The mass spectrum of a given sample typically provides information on protein intensity, mass to charge ratio, and molecular weight. In preferred embodiments of the invention, the molecular weights of proteins in the sample are used as a matching criterion to query a database. The molecular weights are calculated conventionally, e.g., by subtracting the mass of the ionizing proton for singly-charged protonated molecular ions, by multiplying the measured mass-over-charge-ratio by the number of charges for mutliuply-charged ions and subtracting the number of ionizing protons.

10 Various databases are useful in accordance with the present invention. Useful databases include, databases which contain genomic sequences, expressed gene sequences, and/or expressed protein sequences. Preferred databases contain nucleotide sequence-derived molecular masses of proteins present in a known organism, organ, tissue, or cell-type. There are a number of algorithms to identify open reading frames
15 (ORF) and convert nucleotide sequences into protein sequence and molecular weight information. Several publicly accessible databases are available, including, SwissPROT/TrEMBL database which contains substantial entries for a variety of organisms, including, *B. subtilis* and *E. coli*. For other databases, see also, e.g., TIGR Microbial Database, <http://www.tigr.org/tdb/mdb/mdb.html>; VanBogelen et al.,
20 *Escherichia coli and Salmonella: Cellular and Molecular Biology*, ASM Press, Washington, D.C., 1996; <http://pcsf.brcf.med.umich.edu/eco2dbase>; <http://expasy.hcuge.ch/cgi-bin/map2/def?ECOLI.ECO2DBASE>. Information contained in the databases includes, e.g., gene name, protein name, E.C. number, category of function, Swiss-Prot accession code, sequence code for Genbank, Kohara phage location,
25 genetic map location, direction of transcription on the chromosome, predicted molecular weight and isoelectric point from DNA sequence, etc.

One or more databases can be searched using any suitable search algorithm. For example, the SwissProt/TrEMBL database ("Expasy," Swiss Bioinformatics Institute) using the Sequence Retrieval System (SRSWWW) module. In general, any search
30 strategy can be utilized in accordance with the present invention.

Typically, a mass spectrometer is equipped with commercial software that identifies peaks above a certain threshold level, calculates mass, charge, and intensity of

detected ions. Correlating molecular weight with a given output peak can be accomplished directly from the spectral data, i.e., where the charge on an ion is one and the molecular weight is therefore equal to the numerator value minus the mass of the ionizing proton. However, protein ions can be complexed with various counter-ions and adducts, such as Na^+ , and K^+ . In such a case, it would be expected that a given protein ion would exhibit multiple peaks, such as a triplet, representing different ionic states (or species) of the same protein. Thus, it may be necessary to analyze and process spectral data to determine families of peaks arising from the same protein. This analysis can be carried out conventionally, e.g., as described by Mann et al., *anal. Chem.*, 61:1702-1708, 1989.

In matching a molecular mass calculated from a mass spectrometer to a molecular mass predicted from a database, such as a genomic or expressed gene database, post-translation processing may have to be considered. There are various processing events which modify protein structure in a cell, including, proteolytic processing, removal of N-terminal methionine, acetylation, methylation, glycosylation, etc.

A database can be queried for a range of proteins which match the molecular mass of the unknown. The range window can be determined by the accuracy of the instrument, the method by which the sample was prepared, etc. Based on the number of hits (where a hit is match) in the spectrum, the unknown is identified or classified.

A preferred method of the present invention concerns identifying one or more unknown microorganisms in a sample, comprising: searching a sequence database for a plurality of different proteins that have the molecular weights of proteins in a mass spectrum of a sample, wherein said sample comprises a plurality of proteins from one or more unknown microorganisms, whereby said one or more unknown microorganisms are identified.

Identifying is meant in the general sense. For example, when an unknown microorganism is utilized in the aforementioned method, an objective is to determine the character of the unknown. This can mean finding out the particular taxonomic group(s) to which the microorganism belongs, such as its kingdom, phylum, class, order, family, genus, species, variety, and/or strain. By determining that the sample is derived from a bacteria, the sample is thus classified as a bacteria. Identification in this sense can be as precise as the materials and methods allow. For some purposes, it may be enough to

-9-

identify a sample as derived from a set of possible groups; however, other purposes may demand more precision. For instance, it may be enough for certain purposes to describe the sample as comprising a bacteria, as opposed to a protozoa, or a pathogenic bacteria as opposed to a nonpathogenic bacteria.

5 In accordance with the method, a database is searched for proteins which have the molecular weights of protein constituents in the sample. A database is a collection of organized information in a form which can be searched and retrieved by a computer, or other electronic processing means. As described above, the searching can be accomplished usually any suitable, effective, search algorithm that can determine the
10 presence of entries in the database which have the same, or within a specified range, molecular weight of proteins in the mass spectrum of the unknown sample. The database, as mentioned earlier, can comprise genomic sequences, expressed genes, protein sequences, protein molecular weights, etc. If the database contains nucleotide information, then this information can be translated into protein data before the searching
15 step, e.g., by identifying an ORF, proteolytic and cleavage sites, glycosylation sites, methylation sites, and other processing which can influence the mass of a protein. In the case where a DNA database is being used to generate and deduce protein information, the knowledge of the protein's characteristics is indirect. Thus, the searching step can be characterized as searching for proteins which are "predicted" to have molecular
20 weights. A search in accordance with the present invention means, e.g., that a database is queried or probed for the presence of a data which matches or corresponds to the measured data, such as the measured data obtained from a mass spectrum.

According to preferred embodiments of the present invention, the database is search for a plurality of different proteins, i.e., more than one, preferably more than 5,
25 etc. In general, identification reliability will depend on a number of factors, including the number of peaks in a matched spectrum matched to proteins in a database, the number and accuracy of proteins predicted from the genome sequence in the mass range under study, etc. By the term "different," it is meant that the proteins arise from different genes, such as a gene coding for a protease and a gene coding for an amylase.

30 A search strategy can use the information generated by MS, or any other method, to search a database. A simple search and find strategy can be used where the database is queried for proteins which match the molecular weight of the inputted data.

-10-

Fig. 6 is an example of a process of identifying an unknown organism, cell, or other biological material. One or more of the steps depicted in the flow chart can be used to identify an organism in accordance with the present invention. In this example, a mass spectrum of a sample comprising proteins from an unknown organism has already been generated using MALDI/TOF spectrometry. The output from the mass spectrometer is represented as a series of m/z values, where m is the mass of a protein plus the mass of a proton or other charging species, and z is the net number of charges carried by the ion and is used as the initial input 1. The input masses are processed 2, e.g., by subtracting one dalton to correct for the proton added to the protein when it is ionized through gas-phase proton transfer reaction of MALDI. The input data can additionally be processed by determining an average molecular weight or a monoisotopic mass. A protein will typically be represented in a mass spectrum by more than one peak because of the presence in it of more than one isotope. Carbon, for instance, occurs in nature as C-12 or C-13 in a ratio of about 100:1. Therefore, if a compound contains a single carbon, it would be expected that 99% of it would be C-12 and 1% of it would be C-13. The mass spectrum of such compound would therefore have at least two peaks, each corresponding to a different carbon isotope. As the number of carbon atoms in a molecule increases, there is an increasing number of polyisotopic molecules, comprising varying ratios of the different carbon isotopes. A mass spectrum of such a compound would contain multiple peaks for each polyisotopic molecule. A compound containing a plurality of atoms represented by more than one isotope will have a complex pattern of spectral peaks. Such complex spectral information can be processed in a number of ways. A average mass can be calculated, e.g., using the empirical spectral information. Alternatively, a monoisotopic mass can be calculated where a mass is derived where the mass of only one isotope of each atom is represented in the molecule.

Before database searching is initiated, a mass window is set 3 to define a mass range in which matches in the database will be scored as hits. The mass window for a particular query can be set based on various criteria. One consideration is the accuracy of the instrument. For instance, if the instrument can only measure values within three daltons, then the mass window could be for ± 3 Da. Other considerations, include, post-translational processing. The accuracy of the instrument can be determined routinely,

-11-

e.g., using known standards and calibrating the instrument using an external and internal standard.

The processed data resulting from 2 is used as input data to initiate a search 4 of a database containing protein masses. In preferred embodiments, the database comprises nucleotide sequence information which has been analyzed to predict the occurrence of open reading frames (ORF) and the calculated molecular masses of such ORFs. As mentioned above, various public and private databases are available that contain calculated protein mass information, or which can be mined by available software to derive such information. The search mode queries the database for proteins having molecular masses which match up with the molecular masses in the mass spectrum input data 2. For each peak in the input data 2, the database is queried and a list is generated of putative database proteins which match it. A match is identified in the database if it possesses the same mass as the peak, or if it is within the range indicated in the mass window 3.

A first list 6 can be generated which reflects the masses and organismic sources for each match. For example, each mass spectral peak of 2 can be associated with a family of proteins, representing proteins of the same molecular mass but from different organisms and proteins within the mass range set in 3.

The data in 6 can optionally be refined 7 by inputting additional data 8, e.g., from fragmented precursor ions of 1 and collecting data of peptide mass, sequence tag information from mass spectra, or other types of downstream information on the constituent proteins. Such data, or orthogonal information, can, e.g., increase certainty that the identification is correct and/or reduce the number of positive hits identified in a search. When a search identifies X possible proteins in the database which match the query by being within the mass window set in 3, a step 8 can be used to reduce the number of possible hits. When the queried database contains amino acid sequence information (deduced or experimentally-derived), sequence information or proteolytic information can be used to determine which specific hit, in the set of hits identified for the specified mass range, corresponds to the data point of interest in the mass spectrum. For example, amino sequence or composition information obtained from a peak of interest can be used to search the set of hits identified as matching the peak of interest to ascertain which hit contains the sequence information. Sequence can information can

-12-

be highly specific, eliminating all other peaks having the same molecular weight from the list generated in 7. Amino acid composition information can also be used a refining tool, although it may be less specific. Any supplemental information on the physical characteristics of a protein can be used to confirm and/or reduce the number of hits identified in a search, including, sequence information as mentioned, cleavage patterns (chemical or enzymatic), isoelectric point, hydrophobicity as deduced from chromatography, immuogenicity, etc.

The data from 6 or 7 can then be scored to generate an output list 10 which lists the possible organisms sources of the mass spectrum. The identified organismic sources can be ranked based on a number of criteria, including, but not limited to, total number of proteins identified as matching an organismic source, orthogonal information obtained in 8, etc. Table 1, for instance shows that *B. subtilis* contains 12/15 or 80% matching peaks and *E. coli* contains 6/15 or 40% matching peaks. If percent match is the sole criteria, *B. subtilis* would be ranked above *E. coli*.

Optionally, the proteins which are unidentified (e.g., the three proteins listed in Table I for *B. subtilis*) in the list can be subjected to further analysis in an iteration step 10. One reason that a matching protein is not identified in the database may be that the protein is subjected to post-translational modifications and therefore does not have the molecular weight predicted by ORF analysis.

An advantage of the present invention is that it can be independent of the specific ionization technique and mass analyzer utilized, alleviating the requirement for rigorous reproducibility, crucial in currently used fingerprint-based approaches. The approach introduced here is independent of relative signal intensities in the mass spectrum. It does not even require that the same set of proteins be expressed and/or detected in each analysis of the same organism, only that a set is characteristic so that it can be associated with a microorganism source. The particular choices of sample preparation, ionization and mass analysis for obtaining mass spectra are not restrictive for the described approach, which also has a potential to be used for identification of cells from individual tissues.

The present invention can be used in variety of different ways and settings and has useful applications in the lab, field, and environmental testing. For example, it can be used in human and veterinary medicine to diagnose normal and pathological

-13-

conditions from biological materials, such as blood, plasma, urine, sperm, fecal matter, and saliva. The present invention is also useful in research and industry. Food samples can be obtained from food materials suspected of contamination.

FOOD-40566

EXAMPLES

For the purpose of illustrating the feasibility of the method MALDI TOF mass spectrometry was employed. The described database search method is not restricted to that specific instrument combination and sample preparation. Sinapinic acid (SA) or α -cyano-4-hydroxycinnamic acid (CHCA) 50 mM in 70:30 CH₃CN:H₂O, and an equimolar mixture of SA and 4-methoxycinnamic acid (MCA) in 70:30 CH₃CN:H₂O were used as matrixes. The microorganisms studied were: *B.subtilis* (strain 168, ATCC# 23857) and *E.coli* (ATCC#11775). They were grown in-house according to standard procedures; 8 g/l nutrient broth (Difco Labs, Detroit, MI) was used as a growth medium, after harvesting the material was centrifuged for 10 min at 10⁴g and washed with water three times prior to lyophilization for prolonged storage at -10⁰ C. Lyophilized vegetative cells were suspended in a 70:30 solution of CH₃CN: 0.1% trifluoroacetic acid at a concentration of 5 mg/ml. *B.subtilis* suspension (0.2 μ l) was deposited on the sample slide before MALDI mass spectrometry. For *E.coli* and *B.subtilis* was prepared by mixing suspensions of the two microorganisms on the slide prior to CPD treatment and MALDI mass spectrometry. In some experiments, an internal mass calibration standard (a solution of bovine insulin and bovine ubiquitin) was added to the *E.coli* sample/matrix mixture on the sample slide in order to increase the accuracy of mass determination. For *B.subtilis*, external calibration of the instrument using a mixture of proteins (bovineinsulin, bovine ubiquitin and horse heart cytochrome C) was performed prior o running the samples. All proteins were obtained from Sigma Chemical Co. (St. Louis, MO).

Positive ion mass spectra (typically from 50 single laser shots rastered uniformly across the sample spot) were recorded in linear mode at 20 kV accelerating voltage and a delay of 0.3 μ s. The estimated N₂ laser fluence was around 10 mJ·cm⁻².

A search by protein molecular mass (M_r) and based on the set of protein molecular weights in the spectra was carried out in the SwissProt/TrEMBL database ("Expasy", Swiss Bioinformatics Institute) using the Sequence Retrieval System (SRSWWW) module at <http://expasy.hcuge.ch/srs5/>. A interactive window ("Alternative Query Form") allows search by a number of classifiers. In case we chose average protein MW as the primary classifier. We selected a ± 3 Da MW window, and the only restriction applied in the query was the choice of the "bacteria" protein subset of the database (in

-15-

earlier release of SwissPROT the identifier "prokaryota" was also available). Thus protein identifies and organismic sources were tentatively assigned for all peaks from the experimental spectra within the range from 4 to 15 kDa.

Under the conditions used, the MALDI spectra (Fig. 2) of *B.subtilis* and *E.coli* contain multiple peaks between 4 and 10 kDa with a signal to noise ratio better than 3. They are listed in Tables 1 and Tables 2, respectively. A database search was performed, based on the observed masses. It was assumed that singly-protonated molecules were detected i.e., a proton mass was subtracted from the observed mass in order to obtain the average M_r . In assigning the respective peaks (i.e. Proteins with M_r within the M_r window chosen: ± 3 Da), the organisms from which each potential protein originates, are also determined. These are presented in Tables 1 and 2. From, Table 1, One microorganism, *B.subtilis*, is identified as the source of 12 of the 15 peaks. There are two "runner-ups" in that example, that provide matches for 6 and 5 of the 15 major peaks. It is evident from Table 2 that 13 *E.coli* proteins match observed peaks (out of 17 total), while one microorganism matches 5 of the 17 peaks. The possibility that unmatched peaks can correspond to alkali cation adducts and/or post-translationally modified products (including proteolytic fragments) of proteins already present in the database will be explored in a software implementation of the described approach.

As already pointed out, there exist inherent problems with the reproducibility of MALDI mass spectra of the same organism - *E.coli* [26,29,30] shows that they do not match each other of the spectrum in Fig. 2b. However, searching the proteome database for masses observed in each spectrum leads to the positive identification of the bacteria in each case (Tables 3-5). This is not surprising since all spectra should reflect the presence of expressed proteins from the same genome. the same type of robustness can be illustrated by comparing the MALDI spectra from the same sample of *E.coli*, obtained in different matrixes. The spectra have different fingerprints - peaks above 5 kDa are more prominent in the spectra obtained with MCA:SA matrix (Fig. 3.a), in comparison to spectra with CHCA matrix (Fig. 2b). However, the database search method results in positive identification of the species in each spectrum (Tables 2 and 7). Effects of incubation time on experimentally obtained mass spectra from *E.coli* have been discussed in the literature [31]. Spectra from *E.coli* harvested after 8 and 32 hours of growth are compared on Fig. 3.b and 2b. Again the overall spectral appearance is

-16-

different for the two samples. Nevertheless the identification is straightforward in both cases (Tables 2 and 7). It appears that experimental factors such as choice of an “appropriate” MADLI matrix, variability in the levels of protein expression, etc. will have limited influence when microorganisms are identified by searching the proteome database.

5

F06080-4405550

References

1. National Center for Biotechnology Information (NIH),
<http://www.ncbi.nlm.nih.gov/Entrez/Gerome/org.html>
2. The *C.elegans* Sequencing Consortium, *Science* **1998**, 282, 2012-2018
- 5 3. Arigioni, F.; Talabot, F.; Peitsch, M.; Edgerton, M.; Meldrum, E.; Allet, E.; Fish, R.;
Jamotte, Th.; Curchod, M.-L.; Loferer, H. *Nat. Biotechnology* **1998**, 16, 851-856.
4. *Bioinformatics: A Practical Guide to the Analysis of Genes and Proteins*; Baxevasis,
A.; Oulette, B., Ed.; Methods of Biochemical Analysis 39; Wiley interscience : New York,
1998.
- 10 5. Roepstorff, P. *Curr. Opin. in Biotechnol.* **1998**, 8, 6-13
6. I. Humphrey-Smith, W. Blackstock. *J. Protein Chem.* **1997**, 231, 1-6.
7. James, P. *Biochem. Biophys. Res. Commun.* **1997**, 231 1-6
8. Kuster, B.; Mann, M. *Curr. Opinions in Struct. Biology* **1998**, 8, 393-400
9. Henzel, W.; Billeci, T.; Stults, J.; Wong, S.; Grimley, C.; Watanabe, C. *Proc. Natl*
15 *Acad. Sci. USA*, **1993**, 90, 5011-5015
10. Mann, M.; Hojrup, P.; Roepstorff, P. *Biol. Mass Spectrum.* 1993, 22, 338-345.
11. Pappen, D.; Hojrup, P.; Bleasby, A. *Current Biology* **1993**, 3, 327-332.
12. James, P.; Quandoni, M.; Carafoli, E.; Gonnet, G. *Biochem. Biophys. Res. Commun.*
1993, 195, 58-64.
- 20 13. Yates III, J.R.; McCormack, A.; Eng, J. *Anal. Chem.* **1996**, 68, 534A-540A.
14. Fenyö, D.; Qin, J.; Chait, B. *Electrophoresis* **1998**, 19, 998-1005.
15. prospector.uscf.edu.
www.proteometrics.com
www.mann.embl-heidelberg.de/Services/PeptideSearch
25 cbrg.inf.ethz.ch/MassSearch.html
expasy.hcuge.ch
www.sequet.diac.uk/mowse.html
16. Jensen, O.; Podtelejnikov, A.; Mann, M. *Rapid Commun. Mass Spectrum.* **1996**, 10,
1371-1378.
- 30 17. Mortz, E.; O'Connor, P. Roepstorff, P.; Kelleher, N.; Wood, T.; McLafferty, F.;
Mann, M. *Proc. Natl. Acad. USA* **1996**, 93, 8264-8267.

18. Yates III, J.R.; Eng, J. US Patent No. 553897 (issued July 23, 1996). "Use of Mass Spectrometry Fragmentation Patterns of peptides to Identify Amino Acid Sequences in Databases."
19. Anhalt, J.P.; Fenselau, C. *Anal. Chem.* **1975**, *47*, 219-225.
- 5 20. Heller, D.; Fenselau, C.; Cotter, R.; Demirev, P.; Olthoff, J.; Honovich, J.; Uy, M.; Tanaka, T.; Kishimoto, Y. *Biochem. Biophys. Res. Commun.* **1987**, *142*, 194-199.
21. *Mass Spectrometry for the Characterization of Microorganisms*; Fenselau, C., Ed.; ACS Symposium Series 541; Am. Chem. Soc.: Washington DC, 1994.
- 10 22. Cain, T.; Lubman, D.; Weber Jr., W. *Rapid Commun. Mass Spectrom.* **1994**, *8*, 1026-1030.
23. Claydon, M.; Davey, S.; Edwards-Jones, V.; Gordon, D. *Nature Biotechnology* **1996**, *14*, 1584-1586.
24. Holland, R.; Wilkes, J.; Rafii, F.; Sutherland, J.; Person, C.; Voorhees, K.; Lay, J. *Rapid Commun. Mass Spectrom.*
- 15 25. Krishnamurthy, T.; Ross, P.; Rajamani, U. *Rapid Commun. Mass Spectrom.* **1996**, *10*, 883-888.
26. Arnold, R.; Reilly, J.; *Commun. Mass Spectrom.* **1998**, *12*, 630-636.
27. Welham, K.; Domin, M.; Scannell, D.; Cohen, E.; Ashton, D.; *Rapid Commun. Mass Spectrom.* **1998**, *12*, 176-180.
- 20 28. Haag, A.; Taylor, S.; Johnston, K.; Cole, R. *J. Mass Spectrom.* **1998**, *33*, 750-756.
29. Wang, Z.; Russon, L.; Li, L.; Roser, D.; Long, S. R. *Rapid Commun. Mass Spectrom.* **1998**, *12*, 456-464.
30. Dai, Y.; Li, L.; Roser, D.; Long, S. R. *Rapid Commun. Mass Spectrom.* **1999**, *13*, 73-78.
- 25 31. Arnold, R.; Reilly, J. *A Study of Bacterial Culture Growth by MALDI-MS of Whole Cells*; Proceedings of the 46th ASMS Conference on Mass Spectrometry and Allied Topics, Orlando, Florida, May 31-June 4, 1998, p. 180.
32. Birmingham, J.; Demirev, P.; Ho, Y-P.; Thomas, J.; Bryden, W.; Fenselau, C. *Rapid Commun. Mass Spectrom.* **1999**, *13*, 604-606.
- 30 33. Das, S.; Yu, L.; Gaitatzes, C.; Roger, R.; Freeman, J.; Blenkowska, J.; Adams, R.M.; Smith, T.F. *Nature* **1997**, *385*, 29-30

-19-

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The preceding preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

5 The entire disclosure of all applications, patents, publications, cited above and in the figures are hereby incorporated in their entirety by reference, including Demirev et al., Anal. Chem., 71:2732-2738, 1999.

10 From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

095044-0001
105000-0001

Table 1. Ranking of organisms according to matched peaks in *B. subtilis* spectrum (Fig. 2.a).

Organism*	Observed mass (Da)												
	3988	4302	4506	4877	4947	4993	5247	5892	6098	6510	6582	6623	6664
<i>B. subtilis</i>	x	x	x	x		x	x	x	x	x		x	x
<i>E. coli</i>		x		x			x	x		x	x		
<i>B. burgdorferi</i>	x	x		x	x			x					
<i>M. tuberculosis</i>							x		x		x		
<i>P. aeruginosa</i>		x				x							
<i>M. leprae</i>										x			x

*Only organisms with more than one matching peak (within ± 3 Da) are listed.

-21-

Table 2. Ranking of organisms according to matched peaks in *E. coli* spectrum (Fig. 2.b).

Organism*	Observed mass (Da)													
	4079	4367	4433	4538	4611	4774	5101	5149	5335	5380	5617	6257	6315	7279
<i>E. coli</i>	x	x	x	x	x		x			x			x	x
<i>H. influenza</i>							x					x		
<i>B. subtilis</i>												x		
<i>M. leprae</i>										x				
<i>B. burgdorferi</i>	x			x		x								
<i>S. typhimurium</i>	x			x				x						
<i>H. pylori</i>									x					
<i>Synechococcus sp.</i>		x												x
<i>M. tuberculosis</i>														x

*Only organisms with more than one matching peak (within ± 3 Da) are listed.

Table 3. Ranking of organisms according to matched peaks in *E. coli* spectrum (Fig. 1.b of Ref. 29).

Organism*	Observed mass (Da)											
	4362	4711	5076	5752	6255	7272	7708	8447	9067	9424	10464	10760
<i>E. coli</i>	x	x	x			x	x	x	x	x	x	x
<i>H. influenza</i>	x		x		x		x	x		x		
<i>B. subtilis</i>					x	x	x	x	x			x
<i>Synechococcus sp.</i>	x					x	x				x	x
<i>H. pyroli</i>		x				x						
<i>M. leprae</i>				x						x	x	x
<i>Rhizobium sp.</i>						x				x		
<i>B. burgdorferi</i>										x		
<i>M. tuberculosis</i>		x		x							x	
<i>S. typhimurium</i>												x

*Only organisms with more than one matching peak (within ± 5 Da) are listed.

-23-

Table 4. Ranking of organisms according to matched peaks in *E. coli* spectrum (Fig. 1.a of Ref. 30).

Organism*	Observed mass (Da)										
	3636	4365	4532	4769	6547	7271	7333	9061	9535	9737	13093
<i>E. coli</i>		x	x			x	x	x	x	x	x
<i>B. subtilis</i>				x		x	x	x	x	x	x
<i>Synechococcus sp.</i>		x	x			x			x	x	x
<i>B. burgdorferi</i>	x		x				x		x		
<i>H. influenza</i>		x								x	
<i>Rhizobium sp.</i>					x					x	
<i>H. pylori</i>				x		x					
<i>M. tuberculosis</i>											
<i>S. typhimurium</i>	x		x							x	

*Only organisms with more than one matching peak (within ± 5 Da) are listed.

Table 5. Ranking of organisms according to matched peaks in *E. coli* spectrum (Fig. 4 of Ref. 26).

Organism*	Observed mass (Da)						
	5100	5380	7280	8320	9070	9530	9740
<i>E. coli</i>	x	x	x	x	x	x	x
<i>B. subtilis</i>					x	x	x
<i>M. tuberculosis</i>					x	x	x
<i>H. pylori</i>			x	x			
<i>B. burgdorferi</i>					x	x	
<i>H. influenza</i>	x				x		
<i>E. cloacae</i>	x		x				
<i>Synechocystis sp.</i>					x		x

*Only organisms with more than one matching peak (within ± 5 Da) are listed.

Table 6. Ranking of organisms according to matched peaks in *E. coli* spectrum (Fig. 3.a).

Organism*	Observed mass (Da)								
	4079	4184	4367	4612	4774	5380	7279	9235	9537
<i>E. coli</i>	x		x	x		x	x	x	x
<i>B. burgdorferi</i>	x	x			x				
<i>M. leprae</i>						x	x	x	
<i>Synechococcus sp.</i>		x	x						
<i>S. typhimurium</i>	x							x	

*Only organisms with more than one matching peak (within ± 3 Da) are listed.

Table 7. Ranking of organisms according to matched peaks in *E. coli* spectrum (Fig. 3.b).

Organism*	Observed mass (Da)									
	4079	4367	4433	4611	4774	5380	6257	6315	6412	9536
<i>E. coli</i>	x	x	x	x		x		x	x	x
<i>B. subtilis</i>							x	x		x
<i>H. influenza</i>							x	x		
<i>B. burgdorferi</i>	x				x					
<i>M. leprae</i>						x			x	
<i>Synechocystis sp.</i>		x							x	

*Only organisms with more than one matching peak (within ± 3 Da) are listed.

-27-

Table 8. Ranking of organisms according to matched peaks in spectrum of *B. subtilis* and *E. coli* mixture (Fig. 5).

Organism*	Observed mass (Da)													
	4611	4774	4877	4963	5013	6098	6412	6510	7203	7279	7334	7724	9235	9536
<i>E. coli</i>	x		x	x			x	x	x	x	x		x	x
<i>B. subtilis</i>			x		x	x		x	x			x		x
<i>M. leprae</i>							x	x	x				x	
<i>Synechocystis</i> sp.							x		x	x				
<i>B. burgdorferi</i>		x	x								x			
<i>H. influenza</i>				x									x	
<i>M. tuberculosis</i>						x						x		
<i>S. typhimurium</i>			x										x	

*Only organisms with more than one matching peak (within ± 3 Da) are listed.

-28-

TABLE 9

Genome	Publication
<i>Haemophilus influenzae</i> Rd	Fleischmann <i>et al.</i> , <i>Science</i> 269 :496-512 (1995)
<i>Mycoplasma genitalium</i>	Fraser <i>et al.</i> , <i>Science</i> 270 :397-403 (1995)
<i>Methanococcus jannaschii</i>	Bult <i>et al.</i> , <i>Science</i> 273 :1058-1073 (1996)
<i>Synechocystis</i> sp.	Kaneko <i>et al.</i> , <i>DNA Res.</i> 3 :109-136 (1996)
<i>Mycoplasma pneumoniae</i>	Himmelreich <i>et al.</i> , <i>Nuc. Acid Res.</i> 24 :4420-4449 (1996)
<i>Saccharomyces cerevisiae</i>	Goffeau <i>et al.</i> , <i>Nature</i> 387 (Suppl.) 5-105 (1997)
<i>Helicobacter pylori</i>	Tomb <i>et al.</i> , <i>Nature</i> 388 :539-547 (1997)
<i>Escherichia coli</i>	Blattner <i>et al.</i> , <i>Science</i> 277 :1453-1474 (1997)
<i>Methanobacterium thermoautotrophicum</i>	Smith <i>et al.</i> , <i>J. Bacteriology</i> 179 :7135-7155 (1997)
<i>Bacillus subtilis</i>	Kunst <i>et al.</i> , <i>Nature</i> 390 :249-256 (1997)
<i>Archaeoglobus fulgidus</i>	Klenk <i>et al.</i> , <i>Nature</i> 390 :364-370 (1997)
<i>Borrelia burgdorferi</i>	Fraser <i>et al.</i> , <i>Nature</i> 390 :580-586 (1997)
<i>Aquifex aeolicus</i>	Deckert <i>et al.</i> , <i>Nature</i> 392 :353 (1998)
<i>Pyrococcus horikoshii</i>	Kawarabayasi <i>et al.</i> , <i>DNA Research</i> 5 :55-76 (1998)
<i>Mycobacterium tuberculosis</i>	Cole <i>et al.</i> , <i>Nature</i> 393 :537 (1998)
<i>Treponema pallidum</i>	Fraser <i>et al.</i> , <i>Science</i> 281 :375-388 (1998)
<i>Chlamydia trachomatis</i>	Stephens <i>et al.</i> , <i>Science</i> 282 :754-759 (1998)
<i>Plasmodium falciparum</i> Chr2 (isolate 3D7)	Gardner <i>et al.</i> , <i>Science</i> 282 :1126-1132 (1998)
<i>Rickettsia prowazekii</i>	Andersson <i>et al.</i> , <i>Nature</i> 396 :133-140 (1998)
<i>Helicobacter pylori</i>	Alm <i>et al.</i> , <i>Nature</i> 397 :176-180 (1999)
<i>Leishmania major</i> Chr1	Myler <i>et al.</i> , <i>Proc. Natl. Acad. Sci. USA</i> 96 :2902-2906 (1999)
<i>Chlamydia pneumoniae</i>	Kalman <i>et al.</i> , <i>Nat. Genet.</i> 21 :385-389 (1999)
<i>Aeropyrum pernix</i>	Kawarabayasi <i>et al.</i> , <i>DNA Research</i> 6 :83-101 (1999)
<i>Thermotoga maritima</i>	Nelson <i>et al.</i> , <i>Nature</i> 399 :323-329 (1999)


Claims:

1. A method of identifying one or more unknown microorganisms in a sample, comprising:
- 5 searching a sequence database for a plurality of proteins that are predicted to have the molecular weights of proteins in a mass spectrum of a sample, whereby said one or more unknown microorganisms are identified.
- wherein said sample comprises a plurality of proteins from one or more unknown microorganisms, and said database is searched for more than one different protein,
- 10 2. A method of claim 1, wherein the sequence database is a protein sequence database.
3. A method of claim 1, wherein the sequence database is a nucleotide sequence database.
4. A method of claim 1, wherein the mass spectrometry data is MALDI-TOF data.
- 15 5. A method if claim 1, wherein the mass spectrometry data is obtained by electrospray on a time-of-flight, quadrupole, or ion trap mass analyzer.
6. A method of claim 1, wherein the sample comprises chemical or enzymatic digested polypeptide fragments.
7. A method of claim 1, further comprising:
- 20 performing a mass spectral analysis on a sample comprising one or more microorganisms.

8. A method of claim 1, further comprising:
identifying molecular weights of proteins in a mass spectrum of said sample.
9. A method of claim 1, wherein said sample comprises at least two different species of microorganisms.
10. A method of claim 1, wherein the sequence database is the NCBI/SwissProt/EMBL database.
11. A method of claim 1, further comprising chemical or enzymatic digestion of a protein in said sample.


09/856044

FIG. 1



09/856044


FIG. 2A



09/856044

T0E080"44095860


FIG. 2B



09/856044

T06080-4409560


FIG. 3A



09/856044

T02080-44095850


FIG. 3B



09/856044

T06080-1105500


FIG. 4



09/856044

FOE030-44095360

FIG. 5



09/856044

**AN EXAMPLE OF A SOFTWARE FLOW CHART FOR
MICROORGANISM AND CELL IDENTIFICATION
BY MASS SPECTROMETRY AND DATA BASE SEARCH**




FIG. 6

DECLARATION FOR PATENT APPLICATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

METHODS FOR IDENTIFYING AND CLASSIFYING ORGANISMS BY MASS SPECTROMETRY AND DATABASE SEARCHING

the specification of which

☐ is attached hereto

☒ was filed on 17 NOVEMBER 1999 as United States Application Number or PCT International Application Number PCT/US99/27191 and (if applicable) was amended on _____

I hereby authorize our attorneys to insert the serial number assigned to this application.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or §365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

PRIOR FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 USC §119			
APPLICATION NO.	COUNTRY	DAY/MONTH/YEAR FILED	PRIORITY CLAIMED

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below.

PROVISIONAL APPLICATION(S) UNDER 35 U.S.C. §119(e)	
APPLICATION NUMBER	FILING DATE
60/108,696	17 NOVEMBER 1998
60/120,679	19 FEBRUARY 1999

I hereby claim the benefit under 35 U.S.C. §120 of any United States application, or §365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. §112.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR §1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

PRIOR U.S./PCT INTERNATIONAL APPLICATION(S) DESIGNATED FOR BENEFIT UNDER 37 U.S.C. §120		
APPLICATION NO.	FILING DATE	STATUS — PATENTED, PENDING, ABANDONED

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected herewith: I. William Millen (19,544); John L. White (17,746); Anthony J. Zelano (27,969); Alan E.J. Branigan (20,565); John R. Moses (24,983); Harry B. Shubin (32,004); Brion P. Heaney (32,542); Richard J. Traverso (30,595); John A. Sopp (33,103); Richard M. Lebovitz (37,067); John H. Thomas (33,460); Catherine M. Joyce (40,668); Nancy J. Axelrod (44,014); James T. Moore (35,619); James E. Ruland (37,432); Jennifer J. Branigan (40,921) and Robert E. McCarthy (46,044)

Send Correspondence to Customer Number 23599



23599

PATENT TRADEMARK OFFICE

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of sole or first inventor (given name, family name)	
Plamen A. DEMIREV	
Signature <i>Plamen A. Demirev</i>	Date 06/13/2001
Residence College Park, Maryland	Citizenship BUL
Post Office Address Department of Chemistry and Biochemistry, University of Maryland, College Park, Maryland 20742 US	
Full Name of additional joint inventor (given name, family name)	
Catherine FENSELEAU	
Signature <i>Catherine Fenseleau</i>	Date June 13, 2001
Residence College Park, Maryland	Citizenship US
Post Office Address Department of Chemistry and Biochemistry, University of Maryland, College Park, Maryland 20742 US	
Full Name of additional joint inventor (given name, family name)	
Signature	Date
Residence	Citizenship
Post Office Address	
Full Name of additional joint inventor (given name, family name)	
Signature	Date
Residence	Citizenship
Post Office Address	
Full Name of additional joint inventor (given name, family name)	
Signature	Date
Residence	Citizenship
Post Office Address	

☐ Additional joint inventors are named on separately numbered sheets attached hereto.